

Cunninghamia

A journal of plant ecology for eastern Australia



Date of Publication:
18/12/2012

ISSN 0727-9620 (print) • ISSN 2200-405X (Online)

The peat-forming spring wetlands of the Strathbogie plateau – floristics and environmental relationships

Fiona Coates and Arn Tolsma

Arthur Rylah Institute for Environmental Research, Department of Sustainability and Environment, 123 Brown Street, Heidelberg, Victoria 3084, AUSTRALIA

Abstract: Spring wetlands on the Strathbogie plateau were mapped using recent aerial photography to estimate their current and former extent. The floristic composition and relationships of the vegetation with topographic, environmental and land use variables were interpreted from an analysis of quadrat data. Wetland condition, threats to their persistence and future management requirements were also identified. More than half of the original wetland vegetation in the study area appears to have been lost, probably as a result of clearing for agriculture. Nine floristic assemblages, identified using agglomerative hierarchical clustering, were identified on acidic to mildly alkaline peat in five different hydrogeomorphic settings. One of these assemblages (Low Open Sedgeland) was uniquely confined to spring-fed mounds, not previously described in Victoria. A key to identification of these groups was developed. Floristic composition was correlated with climate and site disturbance but charcoal throughout sediment cores suggested that historical disturbance regimes included fire. Forested sites at higher elevations where grazing pressure appears to have been less intense were the least disturbed sites. Approximately 60% of the wetlands surveyed were assessed as showing signs of soil moisture loss but there was no evidence that water extraction via dams and known bores was a significant driver of current vegetation composition. Threats requiring management were related to habitat destruction and degradation, dysfunction of physical and biological processes, and changes to disturbance regimes. Establishment of native vegetation buffers and biomass management are likely to be of benefit for future management of spring wetland vegetation.

Key words: spring wetlands, disturbance, grazing, peat, management.

Cunninghamia (2012) 12 (4): 363–383

doi: 10.7751/cunninghamia.2012.12.025

Introduction

Wetlands are a threatened vegetation type worldwide as a result of human impact (Mitsch & Gosselink 2007). Ecosystems which are entirely dependent on groundwater represent less than 1% of the land mass of the Australian continent (Hatton & Evans 1998) and include well known wetlands such as the mound springs of the Great Artesian Basin, the wetlands of the Swan Coastal Plain, karstic ecosystems and riparian red gum stands associated with inland groundwater ecosystems (Hatton & Evans 1998).

Wetlands provide a range of ecosystem services related to catchment hydrology and support distinct plant communities restricted to inundated or seasonally inundated parts of the landscape by virtue of their adaptation to waterlogging (Keddy 2010). In agricultural landscapes, year round water availability has historically made spring wetlands vulnerable to land use pressures (Keddy 2010). Spring wetlands may also be at risk from extractive activities that deplete groundwater reserves because of their value as a year round and permanent water supply.

Classification of wetlands at landscape scale is typically linked to their role as functional units, based on hydrogeomorphic classes (Robertson & Fitzsimons 2004, Mitsch & Gosselink 2007) or hydrological setting (Ramsar Convention Bureau 1991). At the regional or management scale, these classes may be subdivided into ecological units based on the floristic and structural attributes of the vegetation (Fensham *et al.* 2004a).

In Victoria, wetlands have been mapped and classified as 127 different Ecological Vegetation Classes (EVCs), descriptive units encompassing a range of environmental variation from sea level to alpine regions including freshwater and saline systems (Department of Sustainability & Environment 2012). The primary intention of EVC mapping and classification is to provide an inventory of vegetation types for various planning and land management activities (Roberston & Fitzsimons 2004) and condition assessments (DSE 2012).

The delineation of Victorian spring wetland vegetation communities has been difficult, partly because examples are severely altered from their original condition and are patchily distributed across the landscape, and partly owing to a lack of systematic surveys. For example, the current EVC description for 'Spring-soak Woodland' is derived from only a handful of remnants in north-eastern Victoria, almost all of which consist of only a subset of species with fidelity to the EVC. Within any single EVC, a range of vegetation communities also occurs, according to local site environmental variation and/or disturbance history.

The relationships between wetland plant communities and environmental variables remain generally unquantified in south-eastern Australian classifications (Roberston & Fitzsimons 2004). To date, the most comprehensive ecological studies of spring wetland vegetation have focused on mound springs associated with the Great Artesian Basin (Fensham & Fairfax 2003, Fensham & Price 2004, Fensham *et al.* 2004a, Fensham *et al.* 2004b). Other studies have investigated the influence of water regime on vegetation patterns and ecosystem function with a view to formulating management directions at broader scales, although field-based vegetation community level studies are still generally lacking (Eamus *et al.* 2006, Raulings *et al.* 2010).

Spring wetlands are rare in the Victorian landscape, particularly at lower elevations. These wetlands are subject to a range of threats, the majority of which derive from land uses incompatible with their preservation and it is likely that most have been destroyed, or severely altered from their original condition since European settlement. To date, no systematic surveys have been conducted in spring wetlands or related groundwater dependent vegetation in Victoria, other than in alpine and subalpine areas (Ashton & Hargreaves 1983, Wahren *et al.* 1999, Whinam *et al.* 2003, Coates *et al.* 2012). Consequently, their floristic composition and contribution to biodiversity are poorly understood and any prescriptions for management are hampered by a current lack of knowledge of floristics, condition, threats and environmental relationships.

The wetlands surveyed in this study support remnant vegetation communities that have developed on and around groundwater-fed springs on the Strathbogie plateau in central Victoria and represent a unique ecosystem in the State. To contribute to an understanding of their management requirements applicable at the site scale, the aims of the current study were to, 1) describe the floristic variation, structure and environmental correlates of remnant spring wetlands; 2) determine their current and former extent, and 3) determine their current condition and threats to their persistence.

Methods

Study area

The study area covers 2 568 km² (Phillips *et al.* 2002) and is roughly 150 km northeast of Melbourne (37° 00' 30" S, 145° 27' 50" E; Figure 1). The plateau is part of a batholith consisting of two coarse-grained granite plutons, formed from a Middle to Upper Devonian intrusion (370–390 m.y.), with a mildly dissected surface of rolling to hilly tableland at about 320–700 m asl. The plateau straddles the Broken River and Goulburn River catchments and exerts major control over river patterns and groundwater flow (Phillips *et al.* 2002).

The hydrogeology of the Strathbogie Plateau has not been studied in detail but is thought to conform to a conceptual model for fractured rock aquifers, described by Stewardson *et al.* (2009). In their model, fluvial erosion through the weathered batholith surface has established valley floors in the erosion-resistant basement rock. Drainage is to the west-southwest, following the orientation of the basement surface, also the dominant surface flow direction of both plutons. Stewardson *et al.* (2009) proposed that the hills surrounding these valleys consist mostly of isolated fractured rock aquifers. Groundwater is discharged from these aquifers via springs located where river valleys intersect with basement rock. Owing to the orientation of the basement surface, springs on the west-southwest side of hills are likely to have a more persistent flow. More intermittent flows can be expected where groundwater discharge occurs at the intersection of hill surfaces and the aquifer water table. Springs are also most likely to occur along valley margins rather than within drainage lines. Palustrine-type wetlands (Ramsar Convention Bureau 1991) have developed around spring outlets at higher elevations on the plateau.

The nearest weather station is at Strathbogie, roughly 40 km to the north of the study area (36° 51' 50" S, 145° 44' 51" E). January and February are the hottest and driest months, with a mean maximum temperature of 27.3°C, mean minimum of 11.6°C and mean monthly precipitation of 46 mm (Bureau of Meteorology, online). The elevation of the plateau mediates the summer heat of the surrounding plains. Prevailing winds are variable, tending southerly to northerly and reaching the highest speeds (11–13 km/hr on average) between early spring and late summer, weakening over autumn and winter




Fig. 1. Location of the study, shown as the area of Strathbogrie granite.

to about 9 km/hr. Winters are relatively cold and wet. The average maximum temperature in July is 10.1°C, the average minimum is 1.6°C, and mean precipitation is 123 mm. Rain (>1mm) falls for 91 days per year on average and snow may fall occasionally (Bureau of Meteorology, online). Rainfall and temperatures are likely to vary over the plateau with altitudinal and topographic variation.

The pre-European vegetation of the study area generally consisted of “lightly forested grassland, dotted with large granite outcrops and numerous springs and creeks” (McCall 1982 cited in Gubbins 2010). However, depressions where trees were cleared are still clearly visible in paddocks and imply that the vegetation may have been relatively dense on lower slopes. Much of the native vegetation was cleared in the mid to late 19th century and replaced with herbaceous pasture species.

The major current land use is grazing by domestic stock, predominantly cattle but also sheep (Carr *et al.* 2006). The consequences of disturbance by stock include soil compaction, localised turbidity, eutrophication via faeces and urine, pugging of soils and destruction of peat, and ponding of water in micro-topographic relief features resulting in increased water loss by evaporation (Carr *et al.* 2006). Water extraction by bores and construction of farm dams above or below springs are common on freehold land in the study area (Carr *et al.* 2006). Other disturbance includes draining of wetlands via channels and planting of high water-use trees, notably blue gum (*Eucalyptus globulus*) plantations adjacent to or near wetlands. The effect of plantations on watertables

is unknown. Other major impacts to wetland condition are weed invasion by pasture species, inadvertent off-target damage to vegetation by poor herbicide use, and broadacre application of fertilisers (Carr *et al.* 2006).

Bushfires were first recorded in 1851 in the region (White *et al.* 1990), although the precise area or whether the Strathbogrie plateau burnt, is not clear. Fires occurred in the first half of the century in 1904, 1908, 1922, 1939 but none have been recorded since 1983 (Gubbins 2010). Fires are likely to have been extensively used to clear land around the time of settlement between 1841 and the early 20th century (Mackrell 1977, Mc Call 1982, both cited in Gubbins 2010), although clearing still occurred in the 1940s (J. Webb, “Kobyboyne” pers. comm.).

The study sites encompassed a range of peat-forming spring wetlands (mires). Peat was defined as organic sediments formed on permanently saturated organic soils and exceeding 20% dry weight (Whinam & Hope 2005, Hope *et al.* 2011). More specifically, the wetlands consist of fen and swamp forest (or wooded fen), as adapted for usage within Australasia (Whinam & Hope 2005, Hope *et al.* 2011). The term ‘fen’ is used to describe vegetation dominated by peat-forming graminoid species, while ‘swamp forest’ describes vegetation dominated by tall shrubs and trees (Whinam & Hope 2005). Sites allied with baseflow rivers and streams, flood plains or alluvial environments associated with flowing water were excluded.

Mapping

Wetlands in the study area were mapped in GIS (Arcview 3.3) using geo-rectified aerial photography (50 cm pixel resolution) collected in 2007. Mapping was also informed by field notes compiled from inspection of total wetland extent. These notes described the full range of vegetation communities and threats (see below). This allowed us to identify various vegetation types in the aerial images. The average size of the wetlands assessed was approximately 4 ha.

Additional data were extracted from 311 grid locations from a previous survey (Carr *et al.* 2006) and mapped if detectable in the imagery. In the absence of ground-truthing it was not possible to distinguish precise vegetation groups from aerial photographs. Therefore, all wetlands or sections thereof were mapped as one of four broad vegetation types: forest/woodland; shrubland; sedgeland/reedland, and 'remnant' (patches of *Juncus sarophorus*, *Carex* spp. and other non-woody vegetation associated with seepage areas in paddocks). The 'remnant' category was assumed to have once supported wetland vegetation and its total area was included to calculate the estimated (and conservative minimum) loss of wetland vegetation in the study area since European settlement. We made a number of assumptions to identify likely remnants, namely: the Strathbogies were largely forested at European settlement and paddocks are therefore an artefact of clearing, without the wetland vegetation that might once have existed there; peaty soils in paddocks once supported wetland vegetation communities all of which remains is a small number of rushes or sedges, mainly *Juncus sarophorus*; former areas of wetland are visible on aerial photography as darker areas most likely indicating wetter soils.

Vegetation survey

Forty-one wetlands identified in a previous study (Carr *et al.* 2006) were selected according to the following criteria: a range of condition states; accessibility; distinguishable on colour aerial photographs; spanned the geographic and altitudinal ranges of the study area; likely to represent the range of floristic compositional variation and with a known management history.

To describe the floristic variation of the wetlands, percent cover (estimated visually) of all vascular plant species and *Sphagnum* spp. was recorded in quadrats (10m x 10 m) in each of the 41 wetlands in September to November 2008. Quadrats were located near to the point of groundwater discharge and where the vegetation was considered most intact. Plant nomenclature follows Walsh & Stajsic (2006); community structural typology follows Specht *et al.* (1974) and EVC typology follows Department of Sustainability & Environment (2012).

In each quadrat geographic position; quadrat position in the wetland (centre, middle, outer); elevation (metres ASL); slope aspect and angle (using a Suunto compass and clinometer); percent cover (estimated visually to the nearest 10%) and depth of standing water were recorded. Synthetic climate parameters for a range of rainfall and temperature variables

were generated for each quadrat using the BIOCLIM module (Nix and Busby 1986) of the ANUCLIM package (Houlder *et al.* 2005).

A condition descriptor derived from the proportion of three variables was used as a surrogate for site condition in each quadrat [(% pugging + % tracks + % weed cover)/100]. The proportion of the ground covered by pugging and tracks was estimated visually. Weed cover (percent), weed species richness and native species richness were derived from the floristic survey. Other variables recorded were proportion of the wetland perimeter buffered by vegetation with >25% native species cover (estimated in the field and using GIS); overall wetland buffer width (estimated in the field and using GIS); distance and direction (upslope, downslope, across slope) to the nearest dam; distance to the nearest bore, and years fenced.

A sediment core (50 mm x 1 m gouge auger, Dormer Engineering) was taken in each quadrat to determine the depth of the peat layer and the depth to the impermeable layer. The presence of charcoal particles was recorded as an indication as to whether sites were likely to have been burnt at any time in the past. Three soil samples were collected at a depth of 15 cm in each quadrat and bulked for the following analyses: pH of the A horizon (TPS Digital pH meter calibrated with a buffer solution to pH 7, 1:1 ratio of soil to de-ionized distilled water, using 20 gm of soil : 20 ml of water); % moisture and % organic content of the A horizon (measured using loss on ignition, Dean 1974).

To determine the degree to which compositional variation might be associated with landform morphology, each quadrat was assigned to one of five hydrogeomorphic classes based on landform setting and hydrologic dynamics (Semenuk & Semenuk 1997). These classes were derived from slope angle and land surface shape and fell into two categories. Topogenous sites were those situated on more or less flat terrain below slopes or within basins, with no outlet, a single outlet or both inlets and outlets (Ryadin & Jeglum 2006). These sites appeared to receive water from springs situated throughout the drainage line and/or lateral slopes, as well as a significant amount from sub-surface flows and surface runoff. Soligenous sites were generally associated with slope-breaks, receiving groundwater at its point of expression above the surface, with only minor inflows from soil water drainage and surface runoff (Ryadin & Jeglum 2006).

Hydrogeomorphic classes were: 1, Vales: broad depressions flanked by hillslopes <5° (topogenous); 2, gullies: flanked by hillslopes >5° (topogenous); 3, slopes <5°: flat or near flat, or with a distinct concave depression (soligenous); 4, slopes >5°: sloping, or with a distinct concave depression (soligenous); and 5, mounds: convex formations associated with spring vents (soligenous).

Threats

Threats, management issues and other site parameters that might influence wetland condition or function, or might require management in future were noted during field mapping. These included: land use; recent or planned

changes to land use/management; physical threats; vegetation connectivity (as a measure of overall intactness and fragmentation); vegetation zonation (% of identifiable floristic assemblages intact, estimated visually in the field and from aerial photography); the proportion of wetland soils that were dry/damp and wet/waterlogged (estimated in quadrats and data combined for each wetland site) and any obvious qualitative or quantitative changes to the water regime (Department of Sustainability & Environment 2006). Dry soils clearly contained no or very little moisture and damp soils had limited or no ability to release excess water when a handful of soil was manually compressed; saturated or waterlogged soils were identified as having excess water present, which was freely released with or without light compression.

Analytical methods

To identify vegetation groups, floristic data were analysed by hierarchical agglomerative clustering using average linkage (Oksanen 2011). Species that occurred in fewer than 10% of quadrats were excluded. Compositional dissimilarities were calculated using the Bray-Curtis co-efficient (Faith *et al.* 1987). Unique clusters were identified at the nine group level, considered sufficient to represent the range of vegetation communities observed in the field. A dendrogram was constructed to display group similarity and a dichotomous key was devised to assist with recognition of the vegetation groups in the field.

Differences in mean soil organic and moisture content were compared among hydrogeomorphic classes using a One-way Analysis of Variance. Quadrat variables according to floristic groups were compared using interval plots (Minitab 2010). Pearsons Product-Moment correlation co-efficient was used to test for significant relationships between organic and moisture content, pH, slope angle, precipitation, altitude. Results were considered significant if $P < 0.05$.

The Bray Curtis dissimilarity matrix of floristic data was then ordinated using non-metric multidimensional scaling (NMDS) from several random starts until a solution with the smallest acceptable stress was reached. To determine an appropriate dimensionality, the solution was chosen that provided the most reduction in stress (McCune & Grace 2002). Quadrats were then plotted in the ordination space and labelled according to cluster group membership. To determine the degree to which compositional variation could be explained by wetland management and/or environmental attributes, directional vectors for the variables measured

during the quadrat survey were fitted through the ordination space (Kantvilas and Minchin 1989). Only those variables which satisfied the assumption of linearity and normality were used, tested by fitting surfaces of each of the variables to the ordination and using the Johnson transformation (Minitab 2010).

Data manipulation, statistical analyses and graphics were conducted using Program R Version 2.14.1 (R Development Core Team 2011) with packages MASS (Venables & Ripley 2002) and Vegan version 2.1–14 (Oksanen *et al.* 2012) and Minitab® 16.1.0 Statistical Software (Minitab Inc. 2010).

Threats were compiled and allocated to ecological risk categories (Auld & Keith 2008) to give general context to address potential management issues.

Results

Mapping

A total area of 1646 hectares of wetlands vegetation and remnant wetland vegetation was mapped from aerial photos, representing 0.65% of the total study area. Four broad structural vegetation types were recognised at a landscape scale, representing a gradient of canopy cover: (i) treeless vegetation of fens dominated by rushes, reeds or sedges; (ii) shrublands; (iii) swamp forest and woodland and (iv) remnants dominated by rushes and grasses.

‘Remnant’ accounted for most (58%; Table 1) of all mapped wetland vegetation and forest/woodland the least (2.5%; Table 1). Roughly one quarter of the vegetation mapped was dominated by sedges and reeds and roughly 15% was shrub-dominated (Table 1).

The full extent of wetlands could not be accurately assessed from aerial imagery for two reasons. Firstly, wetlands were difficult to detect when they were under tree canopy, and the limited areas mapped within the forest or woodland category were mostly those that had been detected during the field surveys. Therefore, the total area of this wetland type is likely to be substantially underestimated. Secondly, in many parts of the study area, particularly in the south, wetland vegetation that would be expected to occur around the numerous dams and drainage lines was not discernible in aerial photos.

Table 1. Broad wetland vegetation types mapped within the study area (1,646 ha).

Vegetation type	Number of polygons mapped	Mapped area (hectares)	% of mapped wetlands
Forest/woodland	65	41.8	2.5
Shrub-dominated	303	247.7	15.1
Sedge/reed-dominated	572	398.2	24.2
Remnant	805	958.6	58.2

Vegetation survey

Two hundred and three species were recorded from 68 quadrats in 47 wetlands. Quadrats were fairly uniformly spread over a range of altitudes (300–620 m). Fewer quadrats were sampled at lower altitudes (20% were below 500 m) than at higher altitudes, indicating that these sites were relatively rare. The introduced pasture species *Holcus lanatus*, *Anthoxanthum odoratum* and *Lotus corniculatus* were ubiquitous components of ground layers, and were recorded in more than 70% of quadrats.

The majority of quadrats (41) consisted of soligenous systems associated with a break in steep slopes ($>5^\circ$) or gentle slopes ($<5^\circ$) where bedrock or an impermeable layer was at, or close to the surface. Just over half of these sites were located on broadly planar land surfaces, with the remainder in broadly concave depressions. Runoff from these sites sometimes converged into distinct drainage lines.

Six soligenous sites were recorded on ‘mounds’ located on steep or gently sloping hillsides. Spring-fed mounds appear to be an undescribed landform element in Victoria. They were distinctly convex and usually about 2–4 m in diameter. Mounds had a clearly defined vent and are likely to have been formed by localised upwelling of groundwater, resulting in unstable, raised areas of saturated peat. Related, larger landform features have been described elsewhere as mound springs, associated with the Great Artesian Basin (Fensham *et al.* 2004a, b), tumulus (organic mound) springs in Western Australia (Blyth & English 1996), and spring-fen mounds or groundwater mounds in Europe, North America and Japan

(Ryadin & Jeglum 2006, Tomita 2008). Topogenous sites were situated in vales (17) and in gullies (7). These sites received water from inflow and from springs situated along banks, or via mounds situated within the main drainage lines.

Cores were taken from 47 quadrats but high water content prevented successful extraction from the remainder of quadrats, particularly in mounds. The stratigraphy of sites was reasonably consistent throughout the study area. All sites, with few exceptions, consisted of an A horizon between 3 and 80 cm deep comprising reddish-brown peat or (rarely) organic loam, overlying grey to yellow-grey medium clays generally from about 85–100 cm and an impermeable layer situated above weathering bedrock. Mounds tended to contain unconsolidated gravels or fine sands in the region above the impermeable layer, at approximately 2 m depth. An increase in moisture was detected below the impermeable layer at two sites, suggesting the presence of perched water tables.

Charcoal was recorded at the boundary between peat and clay layers and throughout the deeper clay layers but less commonly in the upper peat layer. Soil A horizons were acidic (pH 5) to mildly alkaline (pH 7.9). Most sites were peat ($>70\%$ of sites), with the remainder organic loam (i.e. having $<20\%$ organic content).

Moisture content was highly variable among quadrats, ranging from only 20% at the driest site to 95% at saturated sites. The organic content of the A horizon varied from 6% to 80% among quadrats. The highest accumulation of organic content was on mounds, although the sample size was small (6 sites). However, there was no statistically




Fig. 2. Dendrogram produced from agglomerative hierarchical clustering of floristic data. Clusters are labelled by floristic group membership. The lengths of the vertical lines represent Bray-Cutis dissimilarity.

significant difference in mean moisture content or organic content of the A horizon among hydrogeomorphic classes ($P < 0.05$). There was a strong positive relationship between organic content and moisture content ($r = 0.84$, $P < 0.001$), suggesting that peat was better developed in wetlands that showed less signs of drying out, at least toward the wetland centre where quadrats were typically sampled.

Vegetation groups

Nine floristic assemblages were derived from the floristic survey, although there was a degree of overlap in the composition of some of the vegetation groups (Figure 2). There were some significant differences between various vegetation groups for altitude, soil moisture, organic content, rainfall, temperature, native species richness, weed species richness, weed cover, buffer width and length and years fenced, distance to the nearest bore, percent pugging and disturbance (Table 2). However, no group was consistently different from any other and there was a large amount of variation in the data (Table 2).

Group 1. *Phragmites australis* Reedland

Dominated by tall dense to mid-dense stands (> 50% cover) of *Phragmites australis* with a sparse to moderately dense (5–60% cover) mid-stratum of sedges such as *Carex appressa*, *Carex fascicularis* or *Carex gaudichaudiana* (Appendix 1). The ground layer is predominantly introduced species, particularly *Holcus lanatus* and *Lotus corniculatus*. *Eucalyptus camphora* subsp. *humeana*, *Leptospermum lanigerum* and *Dicksonia antarctica* are recorded as occasional emergents. Mean native species richness (7) in this group is half or less than that recorded in other groups, significantly so in some cases ($P < 0.05$; Table 2).

Distributed across the study area at a range of altitudes (380–590 m) but mainly at sites with relatively low mean annual rainfall (829 mm), this community is confined to saturated, topogenous sites with standing water in the central zone of vales or gullies (mean soil moisture = 77%).

Referable to Tall Marsh (EVC# 821).

Table 2. Mean values (\pm SD) of environmental variables recorded for each vegetation group.

Means that do not share a letter are significantly different.

Group	1	2	3	4	5	6	7	8	9
Peat depth (cm)	14 \pm 4.8 ^A	4 \pm 1.4 ^A	26 \pm 27 ^A	28 \pm 18 ^A	20 \pm 11.3 ^A	36 \pm 17 ^A	21 \pm 13 ^A	18 \pm 25 ^A	14 \pm 3 ^A
pH	7.2 \pm 0.3 ^A	6.6 \pm 0.7 ^A	6.4 \pm 0.4 ^A	6.9 \pm 0.2 ^A	6.5 \pm 0.9 ^A	6.6 \pm 0.5 ^A	6.7 \pm 0.6 ^A	6.7 \pm 0.6 ^A	6.9 \pm 0.1 ^A
% organic	43 \pm 20 ^{AB}	28 \pm 23 ^{AB}	24 \pm 21 ^B	63 \pm 12 ^A	34 \pm 15 ^A	43 \pm 18 ^{AB}	52 \pm 21 ^{AB}	52 \pm 26 ^{AB}	38 \pm 28 ^{AB}
% moisture	77 \pm 15 ^{AB}	60 \pm 22 ^{AB}	53 \pm 22 ^{AB}	88 \pm 5 ^A	64 \pm 21 ^{AB}	75 \pm 23 ^{AB}	80 \pm 14 ^A	84 \pm 8 ^{AB}	73 \pm 10 ^{AB}
Altitude (m)	472 \pm 86 ^C	459 \pm 45 ^C	543 \pm 29 ^{AB}	382 \pm 103 ^C	582 \pm 33 ^A	545 \pm 32 ^{AB}	533 \pm 49 ^{AB}	507 \pm 94 ^{AB}	578 \pm 41 ^{AB}
Mean annual precipitation (mm)	829 \pm 78 ^{AB}	793 \pm 24 ^B	947 \pm 70 ^A	811 \pm 124 ^{AB}	845 \pm 96 ^{AB}	904 \pm 108 ^{AB}	902 \pm 68 ^{AB}	933 \pm 55 ^{AB}	844 \pm 46 ^{AB}
Mean annual temperature (C)	12.1 \pm 0.5 ^A	12.1 \pm 0.2 ^{AB}	11.9 \pm 0.2 ^{AB}	12.3 \pm 0.7 ^A	12 \pm 0.5 ^{AB}	11.8 \pm 0.5 ^{AB}	11.6 \pm 0.2 ^B	11.9 \pm 0.3 ^{AB}	11.8 \pm 0.4 ^{AB}
Native species richness	7.3 \pm 3.9 ^C	9.2 \pm 4.9 ^{BC}	21.3 \pm 7.4 ^A	17.6 \pm 2.3 ^{ABC}	17.1 \pm 6.1 ^{ABC}	19.2 \pm 7.3 ^{AB}	15.3 \pm 6.5 ^{ABC}	15.8 \pm 4.6 ^{ABC}	22 \pm 6.4 ^{AB}
Weed species richness	5.5 \pm 2.4 ^B	6.8 \pm 1.5 ^{AB}	6.9 \pm 1.9 ^{AB}	10 \pm 2.1 ^A	7.4 \pm 2.1 ^{AB}	9 \pm 2.6 ^{AB}	5.7 \pm 2.2 ^B	7.2 \pm 2.8 ^{AB}	5.8 \pm 2.1 ^{AB}
Weed cover (%)	19 \pm 17.2 ^{ABC}	43 \pm 12 ^A	25 \pm 11.1 ^{ABC}	31 \pm 26.8 ^{AB}	7 \pm 6.1 ^C	19 \pm 11.5 ^{ABC}	12 \pm 11.9 ^{BC}	13 \pm 9.8 ^{BC}	3 \pm 2.5 ^{BC}
Buffer width (m)	26 \pm 47 ^{AB}	21 \pm 35 ^{AB}	34 \pm 32 ^{AB}	0 ^B	151 \pm 216 ^A	27 \pm 29 ^{AB}	30 \pm 42 ^B	0 ^B	61 \pm 28 ^{AB}
Buffer length (m)	17 \pm 27 ^{AB}	34 \pm 65 ^{AB}	34 \pm 34 ^{AB}	0 ^B	70 \pm 41 ^A	57 \pm 56 ^{AB}	23 \pm 28 ^B	0 ^B	58 \pm 30 ^{AB}
Years fenced	3.2 \pm 4.5 ^{AB}	1.2 \pm 2.2 ^B	3.1 \pm 3.8 ^{AB}	0.1 \pm 0.2 ^B	2.5 \pm 3.8 ^B	4.2 \pm 3.8 ^{AB}	2.6 \pm 3.5 ^B	0.5 \pm 0.5 ^B	11.3 \pm 12.5 ^A
Distance to bore (m)	3617 \pm 1710 ^{AB}	2304 \pm 1681 ^B	1477 \pm 851 ^B	6440 \pm 1831 ^A	3513 \pm 2262 ^B	920 \pm 550 ^B	2588 \pm 1422 ^B	3640 \pm 2251 ^{AB}	2795 \pm 1344 ^B
Distance to dam upslope (m)	0 ^A	12 \pm 27 ^A	0 ^A	40 \pm 89 ^A	15 \pm 25 ^A	0 ^A	27 \pm 49 ^A	142 \pm 275 ^A	20 \pm 40 ^A
Distance to dam downslope (m)	432 \pm 279 ^A	226 \pm 173 ^A	119 \pm 162 ^A	80 \pm 148 ^A	366 \pm 582 ^A	97 \pm 101 ^A	174 \pm 277 ^A	76 \pm 98 ^A	48 \pm 59 ^A
Distance to dam across slope (m)	0 ^A	0 ^A	23 \pm 41 ^A	44 \pm 67 ^A	9 \pm 25 ^A	13 \pm 33 ^A	54 \pm 162 ^A	0 ^A	10 \pm 20 ^A
% pugging	0.5 \pm 0.5 ^B	0.5 \pm 0.6 ^B	2 \pm 3.5 ^B	35 \pm 40 ^A	1 \pm 1.7 ^B	2 \pm 1.5 ^B	1 \pm 1.4 ^B	3 \pm 2 ^B	2 \pm 2.4 ^B
% tracks	1 \pm 2 ^A	3 \pm 2 ^A	2 \pm 1.9 ^A	6 \pm 13.3 ^A	8 \pm 7.7 ^A	4.8 ^A	5 \pm 4.6 ^A	0 ^A	8 \pm 4 ^A
Disturbance	0.2 \pm 0.2 ^{ABC}	0.5 \pm 0.1 ^{AB}	0.3 \pm 0.1 ^{BC}	0.7 \pm 0.4 ^A	0.2 \pm 0.1 ^C	0.3 \pm 0.1 ^{BC}	0.2 \pm 0.1 ^C	0.2 \pm 0.1 ^C	0.1 \pm 0.1 ^C

Group 2. *Juncus sarophorus* – *Carex appressa* Sedgeland

This community is dominated by *Juncus sarophorus* and *Carex appressa* (30–70% combined cover) and a moderately dense to dense ground layer (30–80% cover) mainly consisting of introduced pasture species such as *Holcus lanatus*, *Anthoxanthum odoratum* and *Lotus corniculatus*. *Eucalyptus camphora*, *Acacia melanoxylon* and *Leptospermum lanigerum* are occasional emergents (Appendix 1).

Juncus sarophorus – *Carex appressa* Sedgeland is common on both topogenous vales across a range of elevations (390–500 m) at sites with relatively low mean annual rainfall (793 mm). It occurs on drier sites than group 1 (Table 2) and is commonly seen as remnant wetland vegetation in paddocks along shallow drainage lines and at the margins of wetlands.

Referable to Wet Verge Sedgeland (EVC# 932).

Group 3. *Baeckea utilis* Shrubland

Dominated by moderately dense stands (> 40% cover) of *Baeckea utilis* to 4 m high and occasionally co-dominated by *Epacris brevifolia* or *Leptospermum continentale* (Appendix 1). The ground layer mainly consists of exotic

grasses and herbs (*Holcus lanatus*, *Anthoxanthum odoratum*, *Lotus corniculatus*) but also with a high level of species richness (Table 2). Co-occurring native species include *Poa helmsii* and other monocot species (e.g. *Baumea rubiginosa*, *Arthropodium milleflorum*, *Luzula meridionalis*). Native herbs are less consistently represented but may include *Gonocarpus micranthus*, *Hydrocotyle* spp., *Drosera peltata*, *Craspedia paludosa* or *Euchiton involucratu*s. *Eucalyptus camphora* may be an occasional emergent.

Confined to the northern Strathbogies at higher elevations (500–575 m), *Baeckea utilis* Shrubland tends to occur as isolated thickets in soligenous situations along breaks in gentle or steeper slopes on shallow soils, at the margins of *Eucalyptus camphora* open forest and as discontinuous patches in cleared paddocks. Soil moisture and organic content values were the lowest recorded during the survey (53% and 24% respectively; Table 2).

Referable to Perched Boggy Shrubland (EVC# 185).

Group 4. *Leptospermum lanigerum* – *Baumea rubiginosa* Open Shrubland

This community occurs on wet sites dominated by *Leptospermum lanigerum* (10–30% cover) to 1.5 m

Key to Groups

- 1. Trees absent or sparse (cover <5%), dominated by shrubs, reeds, rushes or sedges2
- 1. Trees present (cover >5%), or if absent or sparse not dominated by rushes, reeds or sedges (shrubs usually present)3
- 2. Shrubs (*Leptospermum* spp., *Baeckea utilis*) dominant8
- 2. Reeds, rushes or sedges dominant10
- 3. Swamp woodland with ground layer or understorey of shrubs or coral fern (>25%)4
- 3. Swamp woodland with ground layer of sedges and/or other ferns5
- 4. Swamp woodland with ground layer dominated by coral fern GROUP 9
- 4. Swamp woodland with understorey dominated by shrubs8
- 5. Swamp woodland with ground layer of sedges, bracken and *Blechnum* GROUP 5b
- 5. Swamp woodland with ground layer mainly of sedges (>25%)6
- 6. Open swamp forest with a ground layer dominated by *Gahnia* or *Carex*7
- 6. Swamp woodland with a ground layer dominated by *Baumea* GROUP 7a
- 7. Open swamp forest with a ground layer dominated by *Gahnia sieberiana* GROUP 5a
- 7. Open swamp forest with a ground layer dominated by *Carex appressa* GROUP 5c
- 8. Understorey consisting of tall shrub thickets (2 m)9
- 8. Understorey consisting of low shrubs (< 2m) GROUP 4
- 9. Open swamp forest or shrubland dominated by *Leptospermum lanigerum* GROUP 6
- 9. Open swamp forest or shrubland dominated by *Baeckea utilis* GROUP 3
- 10. Tall reedland (> 2m) GROUP 1
- 10. Dominated by rushes or sedges11
- 11. Dominated by rushes (*Juncus*) GROUP 2
- 11. Dominated by sedges12
- 12. Swamp woodland, ground layer dominated by *Baumea* GROUP 7a
- 12. Sedgeland, swamp gum absent13
- 13. *Baumea* spp. >25% cover14
- 13. Small graminoids and forbs dominant (< 25% cover) GROUP 8
- 14. *Gleichenia* and *Baumea* spp. co-dominant GROUP 9
- 14. *Baumea* dominant GROUP 7b

(occasionally taller) and *Baumea rubiginosa* with a high diversity of graminoid and herbaceous species in the ground layer (Appendix 1). These include *Holcus lanatus*, *Eleocharis gracilis*, *Juncus planifolius*, *Phragmites australis*, *Schoenus* spp., *Isolepis* spp., *Triglochin striata*, *Utricularia dichotoma* and *Hypericum japonicum*.

The community is found at significantly lower mean elevation (380 m, $P < 0.001$) than any other group except 1 and 2 (Table 2). It occurs in soligenous settings around spring outlets within vales as well as on gentle or steeper slopes to the south of the study area. Soils had high organic and moisture content (Table 2). Disturbance was also significantly higher than in any other group except 1 and 2 ($P < 0.001$; Table 2). The presence of remnant stands of *Leptospermum lanigerum* suggests that this community may eventually regenerate to Thicket Swamp Forest (see below) in the absence of stock disturbance.

Referable to Perched Boggy Shrubland (EVC# 185) and Swamp Scrub (EVC# 53).

Group 5. *Acacia melanoxylon* Swamp Forest

This group contains three related swamp forest communities dominated by *Acacia melanoxylon* distributed over a soil moisture gradient and distinguished by understorey composition. Different disturbance histories are also likely to have determined the current floristic composition of the understoreys but sites also had low weed cover (Table 2). Quadrats were located on gently sloping terrain and within vales and gullies; however, low moisture content (Table 2) suggests that most sites appear to be drying out.

Referable to Fern Swamp (EVC# 721) and Swampy Woodland (EVC# 937).

Group 5a. *Acacia melanoxylon* – *Gahnia sieberiana* Open Swamp Forest

Dominated by *Acacia melanoxylon* with occasional *Eucalyptus globulus* to 25 m with >25% cover (Appendix 1). The ground layer is dominated by *Gahnia sieberiana* (20–40% cover), a sparse cover of grasses (*Poa helmsii*, *Poa labillardieri*, *Microlaena stipoides*) and small sedge or herb species. *Baumea rubiginosa* is occasionally locally dominant. *Baeckea utilis* may also be present and *Eucalyptus camphora* is an occasional co-dominant.

Open Swamp Forest is confined to steeper slopes and gullies at higher elevations (around 600 m) in the north Strathbogies. Sites are on gully slopes or drainage lines and are topo-soligenous. This community is likely to be a remnant vegetation type formerly more widespread in drainage lines.

Group 5b. *Acacia melanoxylon* – *Eucalyptus camphora* Swamp Woodland

Recorded from a single quadrat within state forest but distinguished from groups 5a and 5c by a higher cover of bracken and lower cover of native herbs and small sedges (Appendix 1). Swamp Woodland consists is dominated by

Acacia melanoxylon and *Eucalyptus camphora* to 25 m (20% cover) with a moderately dense ground layer (50% cover) dominated by *Carex appressa*, *Carex fascicularis*, *Blechnum nudum*, *Pteridium esculentum*, *Poa helmsii* and remnant *Dicksonia antarctica*.

This community was recorded on a soligenous gentle slope in one of the most elevated parts of the study area (595 m).

The quadrat is a rare example of *Eucalyptus camphora* forest unmodified by agricultural practices, although past disturbances are likely to have included selective logging in the surrounding dry sclerophyll forest and higher fire frequencies than in agricultural land. Grass and other graminoid cover is lower overall (20% cf. 40%) and bracken cover higher (10% cf. <1%) than in the previous group.

Group 5c. *Acacia melanoxylon* – *Carex appressa* Open Swamp Forest

At most sites this community consists of *Acacia melanoxylon* (to 20 m) with *Eucalyptus camphora*, *Leptospermum lanigerum* and *Dicksonia antarctica* at some sites (Appendix 1). Canopy cover is generally >10% but may be lower at disturbed sites. The understorey is dominated by a dense cover of *Carex appressa* (70–90%) and minor species *Cyperus lucidus* and/or *Blechnum* spp., with a sparse cover (<5%) of introduced species such as *Holcus lanatus* and *Lotus corniculatus*.

Open Swamp Forest occurred in gullies and vales at higher altitudes across the study area. Sites were structurally modified and reasonably species poor, but had relatively low weed cover and were thus considered to be in moderate to good condition. Two sites had been fenced for five years.

Group 6. *Eucalyptus camphora* – *Leptospermum lanigerum* Thicket Swamp Forest

This group is distinguished by well-developed stands of *Leptospermum lanigerum* (>50% cover, to 10 m high) with *Eucalyptus camphora* or occasionally *Acacia melanoxylon* emergent above the dense canopy, generally sparse (<10% cover) but occasionally with up to 25% cover (Appendix 1). The understorey is generally quite open, and mainly consists of a sparse cover of younger *Leptospermum lanigerum* individuals, and blackberry at some sites. The ground layer is open to moderately dense (15–50% cover) and is made up of sedges, grasses and ferns which may include *Gahnia sieberiana*, *Poa helmsii*, *Blechnum minus*, and a sparse cover of *Baumea rubiginosa* or occasionally *Gleichenia microphylla*. Minor ground layer species include a mix of small sedges and herbs (e.g. *Eleocharis gracilis*, *Isolepis* spp., *Acaena novae-zelandiae*, *Gonocarpus micranthus* and exotics (e.g. *Hypochaeris radicata*, *Lotus corniculatus*, *Holcus lanatus*).

The community appears to be confined to cooler, high rainfall areas (500–570 m) on slopes to the north of the study area.

Referable to Swamp Scrub (EVC# 53), Riparian Scrub (EVC# 191) with elements of Fern Swamp.

Group 7. *Eucalyptus camphora* Swamp Woodland

Nearly a third of quadrats (22) were allocated to this group, which contained a degree of floristic and structural variation among quadrats. However, all quadrats were characterised by a moderately dense to dense cover (>50%) of *Baumea rubiginosa*, or less frequently *Baumea planifolia*.

Referable to Swampy Woodland (EVC# 937).

7a. *Eucalyptus camphora* – *Baumea rubiginosa* Swamp Woodland

This vegetation community is characterised by very wet peat with well-developed organic content (Table 2) and a sparse to moderate cover (5–50%) of trees and/or tall shrubs (5–20 m) consisting of *Eucalyptus camphora*, *Leptospermum lanigerum* or occasionally *Baeckea utilis* at drier sites (Appendix 1). The second stratum consists of a sparse cover (< 25%) of smaller shrubs (< 2m high), typically *Leptospermum continentale* and/or *Baeckea utilis*. *Baumea rubiginosa* forms a dense ground layer (50–100%) and *Phragmites australis* may develop on very wet sites excluded from grazing. Minor ground layer species are *Blechnum minus*, *Hydrocotyle* spp., *Epilobium* spp., *Gonocarpus micranthus* and small graminoids such as *Juncus* spp. and *Eleocharis gracilis*. Introduced exotics were generally less abundant than in group 7b.

Eucalyptus camphora – *Baumea rubiginosa* Swamp Woodland was recorded in the central wetland zone at higher elevations (440 m–590 m). Quadrats were higher in native species richness than group 7b (17 cf. 10), with lower mean cover of weeds (7% cf. 20% cover). Almost all sites were recorded on gentle to steep soligenous slopes and occasionally in vales and gullies.

7b. *Baumea rubiginosa* or *Baumea planifolia* Closed Sedgeland

This community is lacking in trees, but may occasionally include a sparse cover (<10%) of tall shrubs (*Leptospermum continentale* and/or *Baeckea utilis*). It is dominated by tall, moderately dense to dense swards (>50%) of *Baumea rubiginosa* which may grow to 2 m high (Appendix 1). *Baumea planifolia* is dominant at some sites, mostly to the north of the study area. Ground layer species generally include grasses, other small graminoids and herbs such as *Poa helmsii*, *Eleocharis gracilis*, *Juncus* spp., *Epilobium* spp. and occasionally *Sphagnum* spp. Introduced exotic species *Holcus lanatus*, *Hypochaeris radicata* and *Anthoxanthum odoratum* are also prominent in the ground layer.

Baumea rubiginosa/*Baumea planifolia* Closed Sedgeland occurs at higher elevations (420–590 m) in the central, mid and outer wetland zones. Sites were mostly located on gentle to steep soligenous slopes, mounds and occasionally in vales.

Group 8. *Eleocharis gracilis* Low Sedgeland

Low Sedgeland is dominated by *Eleocharis gracilis* and a suite of small sedges and herbs (60–80% combined cover) forming a low (<0.3 m) turf (Appendix 1). Associated species typically include *Hydrocotyle sibthorpioides*, *Epilobium* spp., *Drosera peltata*, *Schoenus maschalinus*, *Schoenus apogon*, *Isolepis subtilissima*, *Isolepis inundata*, *Juncus articulatus*, *Juncus planifolius* and *Lotus corniculatus*. *Holcus lanatus* may be locally dominant at the spring vent.

Sites occur on waterlogged, soligenous spring-fed mounds with high organic content (Table 2). The community is mainly distributed in cooler, high rainfall areas but is occasionally




Fig. 3. Distribution of quadrats within the 3-dimensional ordination space, dimensions 1 and 2. Points are labelled by cluster group membership and vectors of maximum correlation; fitted vectors correlation co-efficients (r2) and statistical significance. ** <0.01; *** < 0.001. Vector lengths are proportional to strength of correlation.

found at lower altitudes (350–570 m). Mounds occur within forest and in open paddocks.

No referable EVC.

Group 9. *Eucalyptus camphora* – *Acacia melanoxylon* – *Gleichenia microphylla* Swamp Woodland

This group consists of species rich sites with an understorey characterised by a dense cover of *Gleichenia microphylla* (40–80%). Other understorey species may include *Baumea rubiginosa* and *Pteridium esculentum*. *Acacia melanoxylon* and/or *Eucalyptus camphora* are the dominant tree species (3–20 m, <25% cover) and *Leptospermum lanigerum* may be locally dominant in the second stratum (Appendix 1). The ground layer consists of a very sparse cover of small sedges, herbs and grasses but *Holcus lanatus* and other exotics commonly recorded elsewhere in the study area are sparse.

Sites are located on slopes in cool, elevated areas (520–620 m). All sites are on gentle to steep soligenous slopes but are showing signs of deterioration, as evidenced by large areas of *Gleichenia* either dead or dying.

Referable to Swampy Woodland (EVC# 937).

NMDS attained a minimum stress of 0.17 in three dimensions. In the plane bounded by dimensions 1 and 2, floristic composition was correlated with variables related to climate and disturbance (Figure 3). The least disturbed sites, with fewer tracks, less pugging and lower weed cover, consisted of forest and woodland which were better buffered by native vegetation, and situated in cooler, more elevated locations than disturbed sites. These were *Acacia melanoxylon* Swamp Forest (Group 5), *Eucalyptus camphora* – *Leptospermum lanigerum* Open Swamp Forest (Group 6) and *Eucalyptus*

camphora – *Acacia melanoxylon* – *Gleichenia microphylla* Swamp Woodland (Group 9).

At the opposite end of this trend were treeless sites with higher mean annual temperatures at lower elevations and sites to the north of the study area with higher levels of disturbance. These groups were *Juncus sarophorus* Sedgeland (Group 2), *Baekkea utilis* Shrubland (Group 3) and *Leptospermum lanigerum* – *Baumea rubiginosa* Open Shrubland (Group 4).

A third compositional trend (dimensions 1 and 3, Figure 4), was correlated with the vector for native species richness. Quadrats with high species richness mainly belonged to Group 6 (*Eucalyptus camphora* – *Leptospermum lanigerum* Open Swamp Forest) and Group 9 (*Eucalyptus camphora* – *Acacia melanoxylon* – *Gleichenia microphylla* Swamp Woodland). However, while the vector was highly significant in relation to site positions on the ordination, the raw data indicated that the relationship between richness and vegetation groups was weaker. There was also a weak correlation between floristic composition and the number of years that some of these sites had been fenced. This suggests that stock exclusion may have positive benefits; however, almost all sites that had been fenced for five or more years were located to the north of the study area on soils with relatively low (< 60%) moisture content, implying that compositional variation may be attributable in part, to establishment of species suited to drier soils.

The relationship between floristic composition and distance upslope to a dam was also weakly significant (Figure 4). While dams were further upslope (> 150 m) of quadrats in Group 1 (*Phragmites* Reedland) and Group 2 (*Juncus* Sedgeland), inspection of the raw data revealed that there was no consistent relationship between this variable and other groups.




Fig. 4. Distribution of quadrats within the 3-dimensional ordination space, dimensions 1 and 3. Points are labelled by cluster group membership and vectors of maximum correlation; fitted vectors correlation co-efficients (r^2) and statistical significance. * < 0.05; *** < 0.001. Vector lengths are proportional to strength of correlation.

Current and potential threats

Habitat destruction and degradation

Changes in land cover: trends in land use change included an increase in the area of farm land converted to hardwood or softwood plantations, viticulture and horticulture (10% of sites visited).

Grazing: only seven quadrats were currently grazed (<10%). Pugging and tracks were the most frequently recorded threat, but severity was generally low with the exception of cleared mound areas or very wet areas. Grazing was also the most common land use within 250 m of wetlands (82%).

Weed invasion: high levels of weed cover were recorded. *Holcus lanatus*, *Lotus corniculatus* and *Hypochaeris radicata* were the three most frequently recorded weeds, occurring in more than 80% of quadrats. *Rubus* spp. (blackberry) were recorded in approximately 50% of wetlands but were rarely abundant, probably because of high investment in control measures by landowners.

Earthworks and extraction: excavation in the form of dams or channels or water extraction via pipes was observed in 30% of the wetlands visited. Landforming (e.g. embankments) was observed less frequently (< 10%) and was generally associated with roadworks or redirection of water (e.g. around culverts).

Dysfunction of physical and biological processes

Reduction in wetland area: 60% of wetlands were assessed as showing signs of drying out, defined as having more than 50% of their area consisting of dry or damp soil (as distinct from saturated or waterlogged soil). This observation was made in another guise by long-term landowners who frequently commented on the extent to which areas of land, previously boggy 20 years ago, were now traversable in a vehicle. At least some vegetation zonation (> 25% intact) was detected at one third of wetlands. However, almost 50% of wetlands assessed showed no sign of zonation, with a small proportion having 10–20% of the presumably original vegetation zones remaining.

Fragmentation: assumed loss of connectivity was observed in 40% of wetland systems, as a result of general land clearing.

Degradation and clearing of buffers: vegetation with a significant component (> 25%) of native species surrounding wetlands was absent from or sparse (< 0.1 ha) in 60% of wetlands. Of the remaining wetlands, only one site (on public land) was completely surrounded by forest and there were few wetlands (< 20% of sites) with buffers > 0.5 ha.

Changes to disturbance regimes

Increased interspecific competition: Sedge cover in excess of 50% was measured in 37% of quadrats and greater than 75% cover was measured in 18% of quadrats (predominantly *Baumea rubiginosa* or *Eleocharis gracilis*).

Fire: charcoal was recorded in 52% of quadrats, indicating that fire occurred historically throughout the study area.

Discussion

Floristic composition

Most vegetation communities were referable to published EVCs with the exception of group 8 (*Elaeocharis gracilis* Low Sedgeland), which was confined to an undescribed landform element (spring-fed mounds) in Victoria. None of the other floristic groups identified by clustering was confined to any particular hydrogeomorphic class.

We did not attempt to describe vegetation zonation. Intact zones were mostly indistinguishable in the field, having been disrupted as a consequence of various disturbances since European settlement, altered hydrological states and subsequent changes to vegetation structure and composition. The lack of clear or consistent delineation between floristic groups on the basis of substrate properties, landscape setting, or other variables is not surprising. Rather, the results describe the floristic assemblages currently present within the landscape, across a range of elevations and degrees of disturbance. The results suggest that hydrogeomorphic classification is unlikely to provide a suitable basis for conservation and management of remnant vegetation because landform setting and hydrologic dynamics alone do not differentiate the range of floristic assemblages present within these wetlands at the scale the study was conducted.

The structural and floristic variation within the range of landform settings and differing management histories across the study area make it virtually impossible to reconstruct the pre European vegetation and there were few if any examples of the presumed original forests. It seems reasonably likely that the extant vegetation is derivative from wet forest, swamp woodlands and shrublands or thickets dominated by *Acacia melanoxylon*, *Eucalyptus camphora*, *Leptospermum lanigerum*, *Leptospermum continentale* and/or *Baeckea utilis*, with a groundlayer consisting mainly of sedges (Cyperaceae) but including a range of forbs and at some sites, *Sphagnum* spp.

Sediment cores showed a distinct zone separating peat and clay layers but which frequently contained charcoal particles throughout, a pattern which was remarkably consistent all over the study area. This may indicate some sudden environmental change such as the arrival of Europeans who used fire to clear vegetation and to provide green feed for sheep (White *et al.* 1990). As a consequence, peat development may be relatively recent, with water tables rising in response to tree removal. Recent palynological work in three wetlands in the southern part of the study area dated these sites between 1,100 years to a mere 100 years old, lending support to the notion that some wetlands in the study area are an artefact of vegetation clearing (Gubbins 2010). The transformation of swamp forest to sedgeland has also been seen after burning in some northern hemisphere wetlands (von Grootjans *et al.* 2005). Evidence of charcoal

is common in swamps (Whinam & Hope 2005) and charcoal was also frequently recorded throughout the deeper clay layers, suggesting that fire has also occurred (either naturally or as a result of Aboriginal burning) over the longer-term.

Current variation in vegetation composition of spring wetlands on the Strathbogie plateau is related to climate, intensity of land use and management. The least disturbed sites (lower weed cover, less soil disturbance) consisted of forested vegetation which was more likely to occur at higher elevations where grazing pressure appears to have been historically less intense, occurred in State Forest or was confined to relatively steep slopes not targeted for clearing. All of these sites were well buffered by native vegetation and most were distant from upslope dams although the vector for the latter variable was relatively weak.

Treeless vegetation (Groups 2, 3, 4 and 8) was the most highly disturbed and was often associated with wetlands in open paddocks where the ground was pugged and the vegetation had been grazed down to a short turf. These sites undoubtedly represent the vegetation which has been most highly modified since European settlement. Nonetheless, native species were well represented in some of these groups, notably groups 3, 4 and 8. This implies a functional shift in the vegetation, whereby an array of minor species has been able to flourish with increased light availability and control of competitors under a regime of grazing.

There was little or no evidence in the data that water extraction via dams and known bores was a significant driver of vegetation composition at the quadrat scale; indeed, there were few bores mapped within 100 m of any study site and only 149 licensed bores across the Strathbogie plateau, equivalent to a bore density of approximately 1 per 6 km², estimated as less than 5% of a base-level groundwater discharge rate (Stewardson *et al.* 2009).

Soil properties, including moisture content, were not related to variation in floristic composition and were not significantly correlated with disturbance. This tends to suggest that land use has affected the vegetation more directly through clearing and reducing structural complexity, than indirectly through hydrological changes. However, sampling was focussed in the wetter, central wetland areas, where the effects of drying are less likely to be detectable than in the outer wetland areas. Outer zones with shallower soils and greater seasonality in moisture levels would be expected to show trends toward drying earlier than central zones, and might be adversely impacted by any decrease in water availability.

A more accurate inventory of the current number and extent of bores and dams, and the total volumes of water extraction including groundwater discharge rates and both the distribution and pumping rates of groundwater bores (Stewardson *et al.* 2009) may provide more information. One site was close to Dropmore Spring, used since 1931 and with a flow of 0.03 L/s; however, no comparable data were available for other springs or bores. Stewardson *et al.* (2009) concluded that on average, the influence of groundwater extractions on spring discharge is likely to be small. Thus, the most likely explanation for soil drying is reduced spring

discharge in recent years primarily as a consequence of the sequence of lower annual rainfall totals experienced in the Strathbogie Ranges over the last decade (Stewardson *et al.* 2009).

We also observed obvious signs of soils drying out at forest/woodland sites which were among some of the least disturbed and were excluded from stock grazing, but had clearly deteriorated in recent years. These sites were situated on slope breaks with ground layers characterised mainly by large dead or dying colonies of *Gleichenia microphylla* or by dense stands of *Gahnia sieberiana* and at one site, remnant *Sphagnum*. However, it is difficult to determine to what extent loss of soil moisture and wetland contraction is attributable to the dam above or to longer-term climatic changes. Nevertheless, these soil drying trends were indicated at the scale of overall site condition. This is further supported by the results of the mapping, which determined (conservatively) nearly a 60% loss of wetland vegetation, as indicated by the extent of 'remnant' vegetation in paddocks.

Threats and management

Mapping revealed that wetlands within the study area are likely to have decreased, conservatively, by nearly two thirds of their original extent, as a result of historical clearing and grazing (and possibly interruptions to hydrology). The main impact on vegetation is likely to be loss of vegetation zonation and replacement of indigenous wetland species by native or introduced dryland species. The wetlands surveyed were vulnerable to a range of threats which could be grouped into three categories: habitat destruction and degradation, dysfunction of physical and biological processes and changes to disturbance regimes.

The major consequences of grazing are eutrophication resulting from pollution by urine and faeces, disturbance and destruction of peat, selective grazing of palatable species, a reduction in the area of soil available for survival and reproduction of native species, and conditions more suitable for establishment of weeds. Ubiquitous exotic pasture species have virtually replaced the ground layer at many sites, favoured by high soil moisture, soil disturbance and increased nutrient levels from broadacre fertilising.

The specific measurement of seasonal inundation was beyond the scope of the study, although many landowners drew attention to this as a likely threat. Loss of the natural range of water regimes (extent and degree of saturation) will further exacerbate the loss of vegetation zonation or loss of understorey dominants which characterises wetland systems, and which has already occurred at most sites. This may result from a long-term decrease in precipitation, from human-induced changes in the surrounding landscape, or from both of these factors.

The role of native vegetation buffers is likely to be most beneficial in maintaining connectivity, reducing access to stock, improving water quality by filtering surface run-off, and making an important contribution to local biodiversity. Buffers consisted of forested vegetation as well as native grassland which had established when wetlands had been

fenced off to exclude stock. Some were dominated by an array of native species, in particular *Themeda australis*, but this depended on past land management such as fertilizer use and extent of ploughing. These buffers require management in the form of sheep grazing or preferably burning, to maintain the somewhat serendipitous establishment of native vegetation that has occurred around wetland sites.

An emerging industry on the Strathbogie plateau is conversion of farmland to blue gum plantations. The replacement of herbaceous vegetation with monocultures may have potential to disrupt wetland hydrology and some biological interactions. We speculate that impacts might include a loss of pollinators, an increase in predators and herbivores, loss of resilience to disease and pathogens (Auld & Keith 2008), or a reduction in ground water availability through transpiration.

Recent conservation efforts have focussed on fencing wetlands to exclude livestock from some of the wetlands, almost all of which occur on private property. An unforeseen consequence of exclusion has been an increase in dominance of large, highly competitive reeds and sedges (e.g. *Phragmites australis*, *Baumea rubiginosa*). Ongoing risks to vegetation quality include increased interspecific competition leading to exclusion of other species and reduction in species diversity, particularly minor or rare inter-tussock species such as *Eriocaulon scariosum* (Department of Sustainability and Environment 2005). Advantages at these sites include provision of habitat protection from predators for native fauna by dense ground layers.

Recent experimental trials have found that regular biomass removal by grazing, burning or slashing is likely to be necessary to control large sedges and weeds but restoration of species richness in Closed Sedgeland presents a substantial challenge (Coates & Tolsma 2012). Most wetlands are highly modified and it will be futile to attempt to restore them to some idealised, pre-European state. Ongoing experimental trials over timeframes of three to five years are still needed to determine specific management prescriptions.

Acknowledgements

We thank Simon Casanelia (GBCMA) for project management and support and members of the project Steering Committee for their contributions to project design and for providing additional site records. We also thank the many landowners who allowed us access to their properties and provided valuable and interesting information about the management histories of sites. A number of our colleagues generously assisted with plant identifications: Doug Froid (Pathways Bushland and Environment), Neville Walsh, Niels Klazenga and Val Stajsic (Royal Botanic Gardens, Melbourne). Doug Froid also provided comment on the vegetation classification and assignment of vegetation communities to EVCs, and Kay Morris (ARI) provided comment on the manuscript.

References

- Ashton DH & Hargreaves GR (1983). Dynamics of sub-alpine vegetation at Echo Flat, Lake Mountain, Victoria. *Proceedings of the Ecological Society of Australia* 12: 35–60.
- Auld TD & Keith DA (2008). Dealing with threats: Integrating science and management. *Ecological Management and Restoration* 10 Supplement 1: 79–87.
- Blyth J & English V (1996). Endangered – Tumulus Springs. *Landscape* 11: 47. Bureau of Meteorology (online) http://www.bom.gov.au/climate/averages/tables/cw_082042.shtml [last visited 05/05/12].
- Carr GW, Moysey ED, Mathews S, Froid D, White M, Griffioen P, Morgan J & Rosengren N (2006). The location of Spring Soak and Peatland wetlands in the Goulburn Broken Catchment – Wetland Implementation Plan Stage. Unpublished report to the Goulburn Broken Catchment Management Authority.
- Coates F & Tolsma A (2012). An investigation of spring wetland management techniques. Unpublished report to the Goulburn Broken Catchment Management Authority, Arthur Rylah Institute for Environmental Research, Heidelberg, Victoria.
- Coates F, Cullen P, Zimmer H & Shannon J (2012). *How snow gum forests and sub-alpine peatlands recover after fire: Black Saturday Victoria 2009 – Natural values fire recovery program*. Department of Sustainability and Environment, Heidelberg, Victoria.
- Dean WE Jr (1974). Determination of carbonate and organic matter in calcareous sediments and sedimentary rocks by loss on ignition: Comparison with other methods. *Journal of Sedimentary Petrology* 44: 242–248.
- Department of Sustainability and Environment (2005) Advisory List of Rare or Threatened Plants in Victoria – 2005. Victorian. Department of Sustainability and Environment, East Melbourne, Victoria.
- Department of Sustainability and Environment (2006). *Index of Wetland Condition. Assessment of wetland vegetation. Update–March 2006* (Department of Sustainability and Environment: Melbourne).
- Department of Sustainability and Environment (2012). A Field Guide to Victorian Wetland Ecological Classes for the Index of Wetland Condition, 2nd edition. Arthur Rylah Institute for Environmental Research, Department of Sustainability and Environment, Heidelberg, Victoria.
- Eamus D, Froend R, Loomes R., Hose G & Murray B (2006). A functional methodology for determining groundwater regime needed to maintain the health of groundwater-dependent vegetation. *Australian Journal of Botany*, 54: 97–114.
- Faith DP, Minchin PR & Belbin L (1987). Compositional dissimilarity as a robust measure of ecological distance: a theoretical model and computer simulations. *Vegetatio* 69: 57–68.
- Fensham RJ & Fairfax RJ (2003). Spring wetlands of the Great Artesian Basin, Queensland, Australia. *Wetlands Ecology and Management* 11: 343–362.
- Fensham RJ & Price RJ (2004). Ranking spring wetlands in the Great Artesian Basin of Australia using endemicity and isolation of plant species. *Biological Conservation* 119: 41–50.
- Fensham RJ, Fairfax RJ, Pocknee D & Kelley J (2004a). Vegetation patterns of permanent spring wetlands of arid Australia. *Australian Journal of Botany* 52: 719–728.
- Fensham RJ, Fairfax RJ & Sharpe PR (2004b). Spring wetlands in seasonally arid Queensland: floristics, environmental relations, classification and conservation values. *Australian Journal of Botany* 52: 583–595.
- Gubbins W (2010). *Vegetation Change and the Ecological Effects of European settlement: A Palynological and Historical Investigation of the Strathbogie Ranges*, Victoria. BA Honours Thesis, the University of Melbourne.

- Hatton T & Evans R (1998). *Dependence of ecosystems on groundwater and its significance to Australia*. Occasional Paper No. 12/98, Land and Water Resources Research & Development Corporation, Canberra.
- Hope GS, Nanson R & Jones P (2011). The peat-forming bogs and fens of the Snowy Mountains of New South Wales. Office of Heritage and the Environment, New South Wales Government, Queanbeyan.
- Houlder D, Hutchinson M, Nix H & McMahon J (2005). *ANUCLIM Version 5.1* (Centre for Resources and Environmental Studies: The Australian National University, Canberra).
- Kantvilas G & Minchin P (1989). An analysis of epiphytic lichen communities in Tasmanian cool temperate rainforest. *Vegetatio* 84: 99–112.
- Keddy PA (2010). *Wetland Ecology: Principles and Conservation* (Cambridge University Press).
- Mackrell CW (1977). *Strathbogie Centenary: 1977–1977: A History of Strathbogie* (Strathbogie Centenary Committee).
- McCall H (1982). *Highlands 1882–1892* (Wangaratta Audio Visual Branch).
- McCune B & Grace JB (2002). *Analysis of Ecological Communities*. MJM Software Design (Gleneden Beach: Oregon, USA).
- Minitab (2010). *Minitab® Statistical Software Release 16.1.0* (Minitab Inc: State College, Pennsylvania).
- Mitsch WJ & Gosselink JG (2007). *Wetlands* (John Wiley & Sons, Inc).
- Nix HA & Busby J (1986). *BIOCLIM: A bioclimatic analysis and prediction system* (Division of Land and Water Resources, CSIRO: Canberra).
- Oksanen J (2011). *Multivariate Analysis of Ecological Communities in R: Vegan Tutorial*. <http://cc.oulu.fi/~jarioksa/opetus/metodi/vegantutor.pdf> [last visited 05/05/212].
- Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Simpson GL, Solymos P, Stevens MHH & Wagner H (2012). *Vegan: Community Ecology Package*. R package version 2.1–14/r2120. <http://R-Forge.R-project.org/projects/vegan/> [last visited 23/03/12].
- Phillips GN, Ely KS, Hergt JM, Cownley DG, Paul B & Dickins C (2002). 3-dimensional geometry of the Strathbogie batholith. In *Victoria Undercover: Benalla 2002 Conference Proceedings and Field guide: Collaborative Geoscience in Northern Victoria* (Phillips, G.N. and Ely, K.S. eds), CSIRO Publishing, Collingwood. pp. 55–60.
- Raulings EJ, Morris K, Roache MC & Boon PI (2010). The importance of water regimes operating at small spatial scales for the diversity and structure of wetland vegetation. *Freshwater Biology* 55: 701–715.
- Ramsar Convention Bureau (1991). *Proceedings of the 4th Meeting of the Conference of Contracting Parties. Montreux, Switzerland*. (Ramsar Convention Bureau: Gland, Switzerland).
- R Development Core Team (2011). *R: A language and environment for statistical computing*. *R Foundation for Statistical Computing*. <http://www.R-project.org/> [last visited 23/03/12].
- Robertson HA & Fitzsimons JA (2004). Hydrology or floristics? Mapping and classification of wetlands in Victoria, Australia, and implications for conservation planning. *Environmental Management* 34: 499–507.
- Ryadin H & Jeglum J (2006). *The Biology of Peatlands* (Oxford University Press: Oxford, UK).
- Semeniuk V & Semeniuk CA (1997). A geomorphic approach to global classification for natural inland wetlands and rationalisation of the system used by the Ramsar Convention – a discussion. *Wetlands Ecology and Management* 5: 145–158.
- Specht RL, Roe EM & Broughton VH (1974). Conservation of major plant communities in Australia and Papua New Guinea. *Australian Journal of Botany Suppl. Series No. 7*.
- Stewardson M, Western A & Wallis E (2009). The Hydrology of Wetlands in the Strathbogie Ranges. Unpublished report to the Goulburn Broken Catchment Management Authority. Department of Civil and Environmental Engineering, The University of Melbourne.
- Tomita K (2008). Relationships among vegetation, landform, and sediments in spring-fed wetlands on the Owari Hills and Chita Hills, central Japan. *Geographical Review of Japan* 81: 470–490.
- Venables WN & Ripley BD (2002). *Modern Applied Statistics with S*. Fourth Edition (Springer: New York).
- von Grootjans A, Alserda A, Bekker R, Janakova M, Kemmers R, Madaras M, Stanova V, Ripka J, Van Delft B & Wolejko L (2005). In *Mires—from Siberia to Tierra del Fuego* (Steiner, G.M. ed), Stapfia 85, Linz, Austria. pp. 97–116.
- Wahren C-HA, Williams R & Papst WA (1999). Alpine and sub-alpine wetland vegetation on the Bogong High Plains, south-eastern Australia. *Australian Journal of Botany* 47: 165–88.
- Walsh NG & Stajsic V (2007). *A Census of the Vascular Plants of Victoria*. Eighth Edition (Royal Botanic Gardens: Melbourne).
- Whinam J & Hope G (2005). The Peatlands of the Australasian Region. In *Mires—from Siberia to Tierra del Fuego* (Steiner, G.M. ed), Stapfia 85: Linz, Austria. pp. 397–434.
- Whinam J, Chilcott NM & Morgan J (2003). Floristic composition and environmental relationships of *Sphagnum*-dominated communities in Victoria. *Cunninghamia* 8: 162–174.
- White LA, Gigliotti F & Cook PD (eds) (1990). *A Reconnaissance Survey of the Catchment of the Middle Reaches of the Goulburn River*. (Land Protection Division: Dept. of Conservation, Forests & Lands, Victoria).
- White M, Murphy AH & Downe J (2006). *National Recovery Plan for the Buxton Gum Eucalyptus crenulata* (Department of Sustainability and Environment: Melbourne).

Manuscript accepted 3 December 2012

Appendix 1. Examples of vegetation groups and distribution of frequent species in quadrats

Cover-abundance classes:

1 = 0.5–1%; 2 = 1–5%; 3 = 5–15%; 4 = 15–25%; 5 = 25–50%; 6 = 50–75%; 7 = 75–100%.

Introduced species are indicated by ‘*’.

Group 1. *Phragmites australis* Reedland



* <i>Anthoxanthum odoratum</i>	010001
<i>Baumea rubiginosa</i>	000211
<i>Blechnum minus</i>	000011
<i>Blechnum nudum</i>	000001
<i>Callitriche stagnalis</i>	000001
<i>Carex appressa</i>	003550
<i>Carex fascicularis</i>	000511
<i>Carex gaudichaudiana</i>	130000
* <i>Cirsium vulgare</i>	110100
<i>Cyperus lucidus</i>	010000
<i>Dicksonia antarctica</i>	000001
<i>Epilobium pallidiflorum</i>	001010
<i>Eucalyptus camphora</i>	000020
<i>Galium australe</i>	101101
<i>Glyceria australis</i>	311011
* <i>Glyceria maxima</i>	003000
* <i>Holcus lanatus</i>	542301
<i>Isolepis inundata</i>	000001
* <i>Juncus bufonius</i>	000001
<i>Juncus planifolius</i>	000001
<i>Juncus sarophorus</i>	320001
<i>Leptospermum lanigerum</i>	000010
* <i>Lotus corniculatus</i>	031111
* <i>Mimulus moschatus</i>	000001
<i>Phragmites australis</i>	657556
<i>Poa helmsii</i>	000100
<i>Pteridium esculentum</i>	001001
* <i>Rubus fruticosus</i>	001001
* <i>Rumex crispus</i>	110000
<i>Stellaria angustifolia</i>	000001
<i>Epilobium</i> sp.	001001
<i>Isolepis</i> sp.	000001

Group 2. *Juncus sarophorus* – *Carex appressa* Sedgeland



<i>Acacia melanoxylon</i>	00010
<i>Acaena novae-zelandiae</i>	00111
* <i>Anthoxanthum odoratum</i>	32331
* <i>Briza minor</i>	00001
<i>Baumea rubiginosa</i>	02023
<i>Blechnum minus</i>	00010
<i>Callitriche stagnalis</i>	10000
<i>Carex appressa</i>	01353
<i>Carex gaudichaudiana</i>	20000
<i>Centella cordifolia</i>	00001
* <i>Cirsium vulgare</i>	11000
<i>Cyperus lucidus</i>	10012
<i>Eleocharis gracilis</i>	00011
<i>Epilobium billardierianum</i> subsp. <i>hydrophilum</i>	00010
<i>Epilobium pallidiflorum</i>	10300
<i>Eucalyptus camphora</i>	00010
<i>Glyceria australis</i>	10000
<i>Gonocarpus micranthus</i>	00010
* <i>Holcus lanatus</i>	51534
<i>Hydrocotyle sibthorpioides</i>	00001
* <i>Hypochaeris radicata</i>	01112
<i>Isotoma fluviatilis</i>	00101
<i>Juncus planifolius</i>	00011
<i>Juncus sarophorus</i>	55234
<i>Leptospermum lanigerum</i>	00020
* <i>Lotus corniculatus</i>	21121
<i>Luzula meridionalis</i>	00010
* <i>Mimulus moschatus</i>	00010
<i>Phragmites australis</i>	01000
<i>Poa helmsii</i>	02043
<i>Poa labillardieri</i>	00001
<i>Ranunculus glabrifolius</i>	00001
* <i>Rubus fruticosus</i>	00001
* <i>Rumex crispus</i>	10000
<i>Senecio</i> sp.	00100
* <i>Trifolium dubium</i>	00001
<i>Utricularia dichotoma</i>	00100

Group 3. *Baeckea utilis* Shrubland**Group 4. *Leptospermum lanigerum* – *Baumea rubiginosa* Open Shrubland**

<i>Acacia melanoxylon</i>	0100001	<i>*Anthoxanthum odoratum</i>	33210
<i>*Anthoxanthum odoratum</i>	3134333	<i>*Briza minor</i>	11000
<i>Arthropodium milleflorum</i>	1111000	<i>Baumea rubiginosa</i>	21332
<i>Baeckea utilis</i>	5653673	<i>Blechnum minus</i>	01110
<i>Baumea rubiginosa</i>	0222223	<i>Carex gaudichaudiana</i>	11011
<i>Blechnum minus</i>	0000011	<i>Centella cordifolia</i>	10010
<i>Callitriche stagnalis</i>	0011010	<i>*Centaurium erythraea</i>	10001
<i>Carex appressa</i>	0001010	<i>*Cirsium vulgare</i>	01001
<i>Carex gaudichaudiana</i>	1040001	<i>Drosera peltata</i>	10110
<i>Drosera peltata</i>	0111001	<i>Eleocharis gracilis</i>	23143
<i>Eleocharis gracilis</i>	0131015	<i>Eriocaulon scariosum</i>	10110
<i>Epacris breviflora</i>	2010330	<i>Epilobium gunnianum</i>	00100
<i>Eucalyptus camphora</i>	0100002	<i>Epilobium pallidiflorum</i>	01001
<i>Euchiton involucratus</i>	0011011	<i>Eucalyptus camphora</i>	01000
<i>Geranium potentilloides</i>	0001001	<i>Geranium potentilloides</i>	00011
<i>Gleichenia microphylla</i>	0000001	<i>Glyceria australis</i>	11001
<i>Gonocarpus micranthus</i>	1121300	<i>Gonocarpus micranthus</i>	00110
<i>Goodenia elongata</i>	1001000	<i>*Holcus lanatus</i>	34323
<i>*Gratiola peruviana</i>	0010010	<i>Hypericum japonicum</i>	10111
<i>Histiopteris incisa</i>	0000101	<i>*Hypochaeris radicata</i>	31111
<i>*Holcus lanatus</i>	3513322	<i>Isolepis inundata</i>	00110
<i>Hydrocotyle hirta</i>	1100000	<i>*Isolepis levynsiana</i>	10100
<i>Hydrocotyle sibthorpioides</i>	0001011	<i>Isolepis subtilissima</i>	00100
<i>Hypericum japonicum</i>	0011001	<i>*Juncus articulatus</i>	00023
<i>*Hypochaeris radicata</i>	1211110	<i>*Juncus bufonius</i>	00100
<i>Juncus sarophorus</i>	1000011	<i>Juncus planifolius</i>	31122
<i>Juncus sp.</i>	0010011	<i>Juncus sarophorus</i>	11001
<i>Leptospermum continentale</i>	0105000	<i>Leptospermum lanigerum</i>	03333
<i>Leptospermum lanigerum</i>	0000024	<i>*Lotus corniculatus</i>	22111
<i>Lilaeopsis polyantha</i>	0000011	<i>*Mimulus moschatus</i>	10010
<i>*Lotus corniculatus</i>	1111022	<i>Phragmites australis</i>	03014
<i>Luzula meridionalis</i>	1111101	<i>Poa helmsii</i>	00101
<i>Poa helmsii</i>	3200300	<i>Schoenus apogon</i>	41141
<i>Poa labillardieri</i>	1000010	<i>Schoenus maschalinus</i>	31111
<i>Ranunculus glabrifolius</i>	1000010	<i>*Sonchus asper</i>	11011
<i>Poa sieberiana</i>	0100000	<i>Utricularia dichotoma</i>	10201
<i>Poa tenera</i>	0201000	<i>Triglochin striata</i>	20312
<i>*Rubus fruticosus</i>	0100201		

**Group 5a. *Acacia melanoxylon* – *Gahnia sieberiana*
Open Swamp Forest****Group 5b. *Acacia melanoxylon* – *Eucalyptus camphora*
Swamp Woodland**

<i>Acacia melanoxylon</i>	254
<i>Acaena novae-zelandiae</i>	101
* <i>Anthoxanthum odoratum</i>	010
<i>Baeckea utilis</i>	100
<i>Baumea planifolia</i>	101
<i>Baumea rubiginosa</i>	100
<i>Blechnum nudum</i>	031
<i>Carex appressa</i>	010
<i>Cassinia longifolia</i>	101
* <i>Cerastium glomeratum</i>	011
<i>Chiloglottis vallida</i>	111
* <i>Cirsium vulgare</i>	110
<i>Dichondra repens</i>	010
<i>Eucalyptus camphora</i>	400
<i>Euchiton involucratus</i>	010
<i>Gahnia sieberiana</i>	545
<i>Geranium potentilloides</i>	100
<i>Gonocarpus micranthus</i>	100
<i>Gonocarpus tetragynus</i>	101
<i>Goodenia elongata</i>	100
* <i>Gratiola peruviana</i>	010
* <i>Holcus lanatus</i>	131
<i>Hydrocotyle</i> spp.	111
* <i>Hypochaeris radicata</i>	111
<i>Isotoma fluviatilis</i>	110
<i>Leptospermum continentale</i>	100
<i>Mentha laxiflora</i>	100
<i>Microlaena stipiodes</i>	211
<i>Poa helmsii</i>	100
<i>Poa labillardieri</i>	101
<i>Poa tenera</i>	001
<i>Polystichum proliferum</i>	010
<i>Pteridium esculentum</i>	111
* <i>Rubus laciniatus</i>	101
* <i>Rubus fruticosus</i>	010
<i>Senecio minimus</i>	101
<i>Schoenus maschalinus</i>	010

<i>Acacia melanoxylon</i>	1
* <i>Anthoxanthum odoratum</i>	2
<i>Blechnum nudum</i>	4
<i>Carex appressa</i>	3
<i>Carex fascicularis</i>	2
<i>Cyperus lucidus</i>	1
<i>Dicksonia antarctica</i>	1
<i>Eucalyptus camphora</i>	4
<i>Gahnia sieberiana</i>	1
<i>Galium australe</i>	1
<i>Geranium potentilloides</i>	1
* <i>Holcus lanatus</i>	2
<i>Hydrocotyle hirta</i>	1
* <i>Lotus corniculatus</i>	1
<i>Mentha laxiflora</i>	1
<i>Poa helmsii</i>	2
<i>Pteridium esculentum</i>	3

**Group 5c. *Acacia melanoxylon* – *Carex appressa* Open
Swamp Forest**

<i>Acacia melanoxylon</i>	0524
<i>Acaena novae-zelandiae</i>	1110
* <i>Acetosella vulgaris</i>	0011
* <i>Anthoxanthum odoratum</i>	0101
<i>Baumea rubiginosa</i>	1000
<i>Blechnum minus</i>	1011
<i>Blechnum nudum</i>	1301
<i>Callitriche stagnalis</i>	0011
<i>Carex appressa</i>	6777
<i>Carex fascicularis</i>	1000
* <i>Cerastium glomeratum</i>	0001
* <i>Cirsium vulgare</i>	0111
<i>Cyperus lucidus</i>	0211
<i>Dicksonia antarctica</i>	0130
<i>Eleocharis gracilis</i>	1000
<i>Epilobium billardierianum</i> subsp. <i>billardierianum</i>	1000
<i>Epilobium billardierianum</i> subsp. <i>hydrophilum</i>	1000
<i>Eucalyptus camphora</i>	0500
<i>Galium australe</i>	0121

<i>Geranium potentilloides</i>	0011
<i>Histiopteris incisa</i>	0001
* <i>Holcus lanatus</i>	3111
* <i>Hypochaeris radicata</i>	1011
<i>Isolepis inundata</i>	1010
* <i>Isolepis levynsiana</i>	1000
<i>Isolepis subtilissima</i>	1000
<i>Juncus planifolius</i>	1000
<i>Juncus sarophorus</i>	1000
<i>Juncus sp.</i>	1000
<i>Leptospermum lanigerum</i>	2000
* <i>Lotus corniculatus</i>	2011
<i>Mentha laxiflora</i>	0100
* <i>Mimulus moschatus</i>	0110
<i>Poa helmsii</i>	0100
* <i>Rubus fruticosus</i>	1101
<i>Schoenus maschalinus</i>	1000
* <i>Sonchus asper</i>	0010
<i>Stellaria angustifolia</i>	0010
* <i>Trifolium repens</i>	0010



Group 6. *Eucalyptus camphora* – *Leptospermum lanigerum* Thicket Swamp Forest



<i>Acacia melanoxylon</i>	000120
<i>Acaena novae-zelandiae</i>	111011
* <i>Anthoxanthum odoratum</i>	301004
<i>Baeckea utilis</i>	000110
<i>Baumea rubiginosa</i>	220202
<i>Blechnum minus</i>	130111
<i>Callitriche stagnalis</i>	010101
<i>Carex appressa</i>	100010
<i>Chiloglottis vallida</i>	001100
* <i>Cirsium vulgare</i>	101011
<i>Dicksonia antarctica</i>	000010
<i>Eleocharis gracilis</i>	201212
<i>Epilobium billardierianum subsp. hydrophilum</i>	100002
<i>Eucalyptus camphora</i>	003350
<i>Euchiton involucratum</i>	001111
<i>Gahnia sieberiana</i>	003450
<i>Galium australe</i>	111011
<i>Geranium potentilloides</i>	001011
<i>Gleichenia microphylla</i>	000230
<i>Glyceria australis</i>	001110
* <i>Glyceria maxima</i>	000004
<i>Gonocarpus micranthus</i>	100111
<i>Histiopteris incisa</i>	000110
* <i>Holcus lanatus</i>	312311
<i>Hydrocotyle spp.</i>	001101
<i>Hypericum japonicum</i>	000011
* <i>Hypochaeris radicata</i>	111111
<i>Isolepis spp</i>	010100
<i>Isotoma fluviatilis</i>	001010
* <i>Juncus articulatus</i>	000110
* <i>Juncus bufonius</i>	011000
<i>Juncus planifolius</i>	100101
<i>Leptospermum lanigerum</i>	656656
* <i>Lotus corniculatus</i>	211311
<i>Luzula meridionalis</i>	000011
<i>Mentha laxiflora</i>	000110
<i>Microlaena stipiodes</i>	001110
<i>Poa helmsii</i>	511000
<i>Pterostylis falcata</i>	001010
* <i>Rubus fruticosus</i>	131111
<i>Sphagnum novozelandicum</i>	000100

Group 7a. *Eucalyptus camphora* Swamp Woodland

<i>Eucalyptus camphora</i>	44001223205020
<i>Acacia melanoxylon</i>	01000200001110
<i>Leptospermum continentale</i>	02001000111110
<i>Baeckea utilis</i>	12000000000005
<i>Baumea rubiginosa</i>	55766656555565
<i>Gymnoschoenus sphaerocephalus</i>	00000000050000
<i>Acaena novae-zelandiae</i>	01110110001111
<i>Carex appressa</i>	00000201001201
<i>Carex gaudichaudiana</i>	00001000110000
<i>Gahnia sieberiana</i>	03000200000000
<i>Eleocharis gracilis</i>	00210011211010
<i>Isolepis subtilissima</i>	10000301100100
<i>Juncus planifolius</i>	00101101100100
<i>Juncus sarophorus</i>	00000101102010
<i>Schoenus maschalinus</i>	00000100100100
* <i>Anthoxanthum odoratum</i>	03111011111131

<i>Blechnum minus</i>	00220111111100
<i>Dicksonia antarctica</i>	00000200000000
<i>Phragmites australis</i>	00000232024000
* <i>Holcus lanatus</i>	11111212221221
<i>Poa helmsii</i>	10122121111020
<i>Gleichenia microphylla</i>	50202000430000
<i>Sphagnum novozelandicum</i>	00007000100010
<i>Gonocarpus micranthus</i>	01111010111011
* <i>Hypochaeris radicata</i>	01111111110011
* <i>Lotus corniculatus</i>	00011111111111
<i>Hypericum japonicum</i>	00011001110110
<i>Craspedia paludosa</i>	00001000110000
<i>Eriocaulon scariosum</i>	00100000000000
<i>Epilobium billardierianum</i>	
subsp. <i>billardierianum</i>	00001100010100
<i>Epilobium billardierianum</i>	
subsp. <i>hydrophilum</i>	00110000000000
<i>Epilobium gunnianum</i>	00100010100000
<i>Euchiton involucratus</i>	01000000100011
<i>Galium australe</i>	00000111000100
<i>Geranium potentilloides</i>	00000100101010
* <i>Rubus fruticosus</i>	00000101001110
<i>Goodenia elongata</i>	01000000110000
<i>Hydrocotyle</i> spp.	01011001110001
<i>Centella cordifolia</i>	01010000100001
<i>Luzula meridionalis</i>	00000010010010
* <i>Mimulus moschatatus</i>	00000101001100

<i>Epilobium billardierianum</i>	
subsp. <i>billardierianum</i>	02100001
<i>Epilobium billardierianum</i>	
subsp. <i>hydrophilum</i>	01000000
<i>Epilobium pallidiflorum</i>	10010100
<i>Gahnia sieberiana</i>	00001000
<i>Galium australe</i>	10001000
* <i>Glyceria maxima</i>	00000100
<i>Gonocarpus micranthus</i>	00001110
* <i>Gratiola peruviana</i>	10000100
<i>Hemarthria uncinata</i>	00000001
* <i>Holcus lanatus</i>	13353452
<i>Hydrocotyle</i> sp.	00000010
<i>Hydrocotyle hirta</i>	01000000
<i>Hypericum japonicum</i>	00000010
* <i>Hypochaeris radicata</i>	02111110
<i>Isotoma fluviatilis</i>	10000000
* <i>Juncus articulatus</i>	00000001
<i>Juncus planifolius</i>	00101000
<i>Juncus sarophorus</i>	01121100
<i>Leptospermum continentale</i>	01000133
<i>Lilaeopsis polyantha</i>	00000100
* <i>Lotus corniculatus</i>	11211111
<i>Luzula meridionalis</i>	00000110
<i>Poa helmsii</i>	02013421
<i>Ranunculus glabrifolius</i>	00010100
<i>Poa sieberiana</i>	00000001
<i>Poa tenera</i>	00000001
<i>Pteridium esculentum</i>	02100000
<i>Sphagnum</i> sp.	00001100
<i>Stellaria angustifolia</i>	10010100



Group 7B. *Baumea rubiginosa* or *Baumea planifolia* Closed Sedgeland

<i>Acaena novae-zelandiae</i>	01100000
* <i>Anthoxanthum odoratum</i>	00111142
<i>Baeckea utilis</i>	00003001
<i>Baumea planifolia</i>	00003015
<i>Baumea rubiginosa</i>	77765552
<i>Blechnum minus</i>	00001010
<i>Carex appressa</i>	20010000
<i>Carex fascicularis</i>	00000200
<i>Craspedia paludosa</i>	00000001
<i>Drosera peltata</i>	00000001
<i>Eleocharis gracilis</i>	01101110

Group 8. *Elaeocharis gracilis* Low Sedgeland

* <i>Acetosella vulgaris</i>	01000
* <i>Anthoxanthum odoratum</i>	10110
* <i>Briza minor</i>	10010
<i>Baumea rubiginosa</i>	10011
<i>Carex gaudichaudiana</i>	10000
<i>Dichondra repens</i>	00100
<i>Drosera peltata</i>	00120
<i>Eleocharis gracilis</i>	57644
<i>Eriocaulon scariosum</i>	00010

<i>Epilobium billardierianum</i> subsp. <i>billardierianum</i>	00001	<i>Carex appressa</i>	0010
<i>Epilobium billardierianum</i> subsp. <i>hydrophilum</i>	11010	<i>Centella cordifolia</i>	0100
<i>Epilobium pallidiflorum</i>	10000	<i>Dicksonia antarctica</i>	0012
<i>Eucalyptus camphora</i>	01000	<i>Eleocharis gracilis</i>	0122
<i>Euchiton involucratus</i>	11010	<i>Eriocaulon scariosum</i>	0100
<i>Gonocarpus micranthus</i>	00010	<i>Eucalyptus camphora</i>	1401
<i>Goodenia elongata</i>	00100	<i>Euchiton involucratus</i>	0011
<i>Hemarthria uncinata</i>	01100	<i>Gahnia sieberiana</i>	0104
* <i>Holcus lanatus</i>	12222	<i>Gleichenia microphylla</i>	7674
<i>Hydrocotyle sibthorpioides</i>	30131	<i>Gonocarpus micranthus</i>	1114
* <i>Hypochaeris radicata</i>	11110	<i>Goodenia elongata</i>	0001
<i>Isolepis inundata</i>	01101	* <i>Gratiola peruviana</i>	0011
* <i>Isolepis levynsiana</i>	01010	* <i>Holcus lanatus</i>	1111
<i>Isolepis subtilissima</i>	32000	<i>Hydrocotyle hirta</i>	0001
<i>Isotoma fluviatilis</i>	10000	<i>Hydrocotyle sibthorpioides</i>	0010
* <i>Juncus articulatus</i>	10101	<i>Hypericum japonicum</i>	0111
<i>Juncus bufonius</i>	01010	* <i>Hypochaeris radicata</i>	1111
<i>Juncus planifolius</i>	11111	<i>Isolepis inundata</i>	0101
<i>Juncus sarophorus</i>	01110	<i>Isolepis subtilissima</i>	0111
<i>Juncus</i> sp.	02020	<i>Isotoma fluviatilis</i>	0001
<i>Lilaeopsis polyantha</i>	10000	<i>Juncus planifolius</i>	1111
* <i>Lotus corniculatus</i>	01033	<i>Leptospermum lanigerum</i>	0224
<i>Luzula meridionalis</i>	10010	* <i>Lotus corniculatus</i>	0011
<i>Phragmites australis</i>	20000	<i>Mentha laxiflora</i>	0011
<i>Poa helmsii</i>	00110	<i>Pteridium esculentum</i>	1111
<i>Schoenus apogon</i>	02110	<i>Schoenus maschalinus</i>	0210
<i>Schoenus maschalinus</i>	01332	<i>Utricularia dichotoma</i>	0100
<i>Stellaria angustifolia</i>	00212	<i>Viola hederacea</i>	0110
<i>Thelymitra</i> sp.	00110		
<i>Utricularia dichotoma</i>	10000		



Group 9. *Eucalyptus camphora* – *Acacia melanoxylon* – *Gleichenia microphylla* Swamp Woodland

<i>Acacia melanoxylon</i>	1121
<i>Acaena novae-zelandiae</i>	0111
* <i>Anthoxanthum odoratum</i>	1112
<i>Baumea planifolia</i>	2000
<i>Baumea rubiginosa</i>	4111
<i>Blechnum minus</i>	1010
<i>Blechnum nudum</i>	0013
<i>Callitriche stagnalis</i>	0011